



International Journal of Advanced Community Medicine

E-ISSN: 2616-3594
P-ISSN: 2616-3586
www.comedjournal.com
IJACM 2023; 6(1): 01-06
Received: 01-10-2022
Accepted: 08-11-2022

Shimaa Mohamed Mohamed Koabar
M.Sc., Department of Public Health and Community Medicine, Faculty of Medicine, Tanta University, Tanta, Egypt

Rania Mostafa El-Salamy
Assistant Professor, Faculty of Medicine, Industrial Medicine and Occupational Health Tanta University, Tanta, Egypt

Ayman Hassan Abd El-Zaher,
Professor of Chest Diseases Tanta University, Faculty of Medicine, Tanta, Egypt

Nadira Mansour Hassan
Professor, Department of Public Health and Community Medicine, Faculty of Medicine, Tanta University, Tanta, Egypt

Corresponding Author:
Shimaa Mohamed Mohamed Koabar
M.Sc., Department of Public Health and Community Medicine, Faculty of Medicine, Tanta University, Tanta, Egypt

Effect of dietary counseling on patients with chronic obstructive pulmonary disease admitted to a tertiary care hospital, Middle Delta, Egypt

Shimaa Mohamed Mohamed Koabar, Rania Mostafa El-Salamy, Ayman Hassan Abd El-Zaher and Nadira Mansour Hassan

DOI: <https://doi.org/10.33545/comed.2023.v6.i1a.249>

Abstract

Objectives: is to assess the impact of dietary advice on malnourished patients with COPD.

Methods: an intervention study conducted at the chest hospital of Tanta university. One hundred patients with stable COPD were chosen.

Patients received dietary counseling once a week for 6 months based on specific nutritional guidelines for patients with COPD.

Results: After 6 months of the intervention, the mean difference of COPD assessment test (CAT) improved from (19.47± 5.759) preintervention to (12.98±5.635) postintervention, both had moderate respiratory impairment as measured by CAT score and the mean difference of Modified Research council (MRC) dyspnea scale improved from (2.70±.689) preintervention to (1.99±.772) postintervention, both had mild score on (MRC) dyspnea scale, with significant difference $p < 0.001$. Health-related quality of life (HRQOL) increased from 4% preintervention to 30% postintervention with significant difference. $p = (0.002)$.

Subjective Global Assessment scores (SGA) improved where there was enhancement and shift in malnutrition from 77% (52% SGA B, 25% SGA C), to only 55% (43% SGA B, 12% SGA C). ($p < 0.001$).

Conclusion: Dietary counseling has a great impact on malnourished COPD patients. It improves nutritional status, resulting in increased functional lung capacity and, as a result, HRQOL.

Keywords: Dietary counseling, chronic obstructive pulmonary disease patients

Introduction

Chronic obstructive pulmonary disease (COPD) refers to a group of lung diseases that include chronic bronchitis, emphysema, and refractory (irreversible) asthma. (Nguyen *et al.* 2015)^[19].

Recently, the prevalence of COPD in Egypt has been reported to be 10%, which is similar to the global prevalence. (Nguyen *et al.* 2019)^[17] COPD is ranked as the fifth leading cause of chronic disability. It is considered as the fourth leading cause of mortality in the United States. (Nguyen *et al.* 2019)^[17] Patients with COPD usually suffer from malnutrition, with losing weight developing a long time, and it is being associated with ill health. (Hoong *et al.* 2017)^[13].

Many issues can cause malnutrition in COPD patients, including a higher metabolism due to increased respiratory work, chronic inflammation, medications, recurrent infections, and oral or teeth problems, which can lead to reduced dietary intake. (Hoong *et al.* 2017, Günay *et al.*, 2013)^[13, 10].

Nutrition interventions have been found to improve functional capacity, and quality of life in COPD patients. (Günay *et al.* 2013, Baldwin *et al.* 2011)^[10, 2] However, there is a scarcity of studies from low- and middle-income economies addressing the effect of dietary counseling interventions on COPD, with the majority of studies coming from high-income countries. As a result, we conducted this study to test the influence of dietary counseling on improving malnourished COPD patients.

Materials and Methods

Study Characteristics: This 6-month prospective intervention study was carried out at the chest department of Tanta University. The study population included patients over the age of 18 with stable COPD who were admitted to the chest wards in Tanta university hospital.

All stable COPD patients admitted to Tanta university chest hospital over 6 months duration, (100) patients with stable COPD were chosen.

Patients were instructed to follow COPD dietary guidelines for 6 months through dietary counselling sessions. They were given dietary advice once a week for six months, based on specially developed written dietary recommendations for patients with COPD. (GOLD. 2017)^[9].

Study inclusion criteria: Patients over the age of 18 with a clinical diagnosis of stable COPD according to guidelines of the Global Initiative for Chronic Obstructive Lung Disease without acute exacerbation in the previous week were eligible for the study. (Gold. 2017)^[9].

Exclusion criteria: Included patients who were experiencing an acute exacerbation, who were receiving systemic steroids, or who had any other chronic illness affecting their nutritional status, such as tuberculosis, lung cancer, or any other malignancy.

Nutrition Intervention Sessions: All participants received dietary counseling and dietary leaflets according to the recent nutritional guidelines of BABEN. (Baben. 2016)^[3] They received individual dietary counseling once a week for 6 months at the Chest department of Tanta university according to their length of stay. All participants were given a leaflet with tips on nourishing drinks and snacks. Following hospital discharge, follow-up meetings were held in the Chest department of Tanta university each week at regular times confirmed by telephone. During the follow-up period, participants were offered continuous support and feedback at each session. Messages were constructed for each patient according to approach relied on the Health Belief Model. (Nguyen *et al.* 2015, Nguyen *et al.* 2019, Baldwin *et al.* 2011)^[19, 17, 2].

Study Tools

Tool 1: Tool 1 consisted of a predesigned questionnaire used for data collection that included the following: sociodemographic data: age, residence, sex, marital status, level of education, occupation, telephone number, and smoking status.

We assessed the participants' nutritional status using the following methods:

Tool 1: Nutritional Screening by Subjective Global Assessment: The Subjective Global Assessment (SGA) (Canadian Malnutrition Task Force 2021)^[4] is a simple bedside method for identifying those who would benefit from nutrition care and diagnosing malnutrition. A clinical evaluation is performed after taking a history of recent intake, weight change, and gastrointestinal symptoms. Patients were categorized into 3 categories "well-nourished (SGA-A), mild/moderately malnourished (SGA-B), or severely malnourished (SGA-C)".

Tool (2): Assessing health related quality of life and functional assessment using:

1) The COPD assessment test (CAT test) Questionnaire is used to assess respiratory health status. Each question is divided into 5 categories ranging from (0-5) /0 indicates total absence, 1 indicates very mild, 2 indicates mild, 3 indicates moderate, 4 indicates severe, and 5 indicates Very severe condition. (CAT Development Steering Group. 2018)^[5].

- Score>5 upper limit of normal

- Score5-10 low impairment to respiratory function
 - Score10-20 moderate impairment to respiratory function
 - Score>20 sever impairment to respiratory function
 - Score30-40 very sever impairment to respiratory function
- 2) Dyspnea was evaluated using the Medical Research Council (MRC) dyspnea scale, a 5-point scale on which a score of (1-normal) indicates that physical activity is not limited by dyspnea and a score of (2-mild), (3,4-moderate), or (5-severe) indicates that the patient is too dyspnea to leave the house. (Medical Research Council. 1986)^[15].
- 3) Combined COPD testing for assessment to HRQoL: (Gold 2017)^[9].
- CAT<10, MRC (0-1) good HRQoL
 - CAT>=10, MRC>=2 poor HRQoL
- 4) Outcome measures included the following:
- SGA
 - CAT SCORE
 - MRC DYSPNEA SCALE
 - HRQOL

Sample Size and Statistical Analysis: We took all the patients over a six-month period and followed them for another six months. Resulting in a total of 100 patients.

We used the paired - sample t for quantitative data and the chi-square test for qualitative data to compare differences in before and after the intervention.

SPSS version 21.0 (IBM Corporation, Armonk, NY, USA) was used for all statistical analyses, p values were calculated using two-sided tests, and statistical significance was defined at p equals 0.05.

Results

Table 1 highlights the sociodemographic characteristics of the 100 participants enrolled in the study with a mean age of 53.14±11.67 years, 99% was male and 51% from rural residence, 50% had secondary education, 34% were manual workers, with average duration of occupation 31 years, 51% of them had enough income, 96% were married with mean children number 3.

Table 2 shows the associated medical conditions and smoking status of the patients being studied, where 64% of them had a past history of chronic disease or operation, 39% had a positive family history of allergy/asthma, 73% admitted to the hospital with COPD acute attacks more than two times, mean number of exacerbations 3.93±1.75 times, with average length of stay 8.43±3.88 days. About three quarters were current smokers (77%), cigarette smoking (74.7%) and heavy smokers (89%). with mean duration (32) years also average number of cigarettes per day were (33). Nearly half of them (50%) were dealing with animals and had pets in their home.

Table 3 revealed the effect of dietary counseling regarding SGA, CAT, MRC DYSPNEA, and HRQOL, where, in terms of the influence of dietary counseling, 44 of the 100 individuals enhanced their nutritional status or SGA score, a significant association was detected among group (p<0.001). Where there was enhancement and shift in malnutrition from 77% (52%SGA B (mild/moderate malnutrition), 25%SGA C (severe malnutrition)), to only 55% (43% SGA B, 12% SGA C).

Regarding MRC Dyspnea Scale, there is improvement and shift in SCORE from 31% mild degree to 39%, from 56% moderate score to 48%, from 9% sever to 7%. Also, increase in normal score from 4% up to 6%.

Regarding CAT SCORE, there is improvement and shift in SCORE from 48% moderate score to 56%, from 42% sever to 30% only, also shift to mild condition from 4% to 10%. The mean difference of COPD assessment test (CAT) improved from (19.47± 5.759) preintervention to (12.98±5.635) postintervention, both had moderate respiratory impairment as measured by CAT score and the mean difference of Modified Research council (MRC)

dyspnea scale improved from (2.70±.689) preintervention to (1.99±.772) postintervention, both had mild score on (MRC) dyspnea scale, with significant difference p (<0.001). There was also improvement in HRQoL from 4% pre intervention to 30% postintervention. There was significant difference in the group under study between pre, and post intervention, p-value (<0.001 for all except HRQoL (p-value=0.002).

Table 1: Baseline Sociodemographic characteristic of COPD patients (N=100)

Variable	Number of participant (220) (100%)	
	Frequency	percent
Age		
30-	42	19.1
40-	65	29.5
50-	49	22.3
60-	43	19.5
70-78	21	9.5
Age (Mean±SD) Range (25-78)	50.73±12.388	
Sex		
Male	218	99.1
Female	2	0.9
Residencez		
Urban	138	62.7
Rural	82	37.3
Patient education		
Illiterate	104	47.3
Read & Write.	9	4.1
Secondary	97	44.1
University&Higher	10	4.5
Patient occupation		
Manual	90	40.9
Technical	31	14.1
Employee	48	21.8
Private Work	37	16.8
Professional	7	3.2
Unemployed	7	3.2
Duration of occupation (Mean±SD)	30.45±11.06	
Family income per month		
Not Enough	51	23.2
Enough	125	56.9
Enough and saving	44	20.0
Marital status		
Single	7	3.2
Married	205	93.2
Widow	8	3.6
Number of children (Mean±SD)	3.15±1.34	

Abbreviation: n (number), and SD (standard deviation)/ 30-** means (30 up to 40) the same for the subsequent values

Table 2: Medical history and smoking status of the studied patients (N = 100)

Variable	Number of participant (220)(100%)	
	Frequency	percent
History of chronic diseases (DM, HTN, Hepatic, Renal)		
Yes	137	62.3
No	83	37.7
Family history of allergy/asthma		
Yes	107	48.6
No	113	51.4
Admission Status		
1 st time	13	5.9
2 nd time	24	10.9
More than two times	183	83.2
Number of exacerbations (Mean ±SD)	3.87±1.392	
Length of stay (Mean ±SD)	8.48±2.940	
Smoking status		
Current Smoker	189	85.9
Ex-Smoker	22	10.0
Nonsmoker	9	4.1
Type of smoking		
Cigarette	166	78.7

Shisha	14	6.6
Both	31	14.7
Duration of smoking in years (Mean±SD)	31.29±10.9	
Number of cigarettes per day (Mean±SD)	31.40±9.548	
Dealing with animals(buffalo-cow-sheep-goat):		
Yes	125	56.8
No	95	43.2
Do you have pets at home(cat-dog)		
Yes	111	50.5
No	109	49.5

Table 3: Effect of dietary counseling on SGA, CAT SCORE, MRC Dyspnea Scale:

Item	Pre intervention N %		Post intervention N %		Significance test p-value
Subjective global assessment					
SGA A	23	23	45	45	Chi -square
SGA B	52	52	43	43	
SGA C	25	25	12	12	P= 0.03*
MRC dyspnea scale					
Normal	4	4	6	6	
Mild	31	31	39	39	Chi-square
Moderate	56	56	48	48	P= 0.02*
Severe	9	9	7	7	
COPD assessment test (CAT)					
Mild	4	4	10	10	
Moderate	48	48	56	56	Chi-square
Sever	42	42	30	30	
Very Severe	6	6	4	4	P= 0.017*
HRQOL					
Poor	96	96	70	70	Chi-square
Good	4	4	30	30	P= 0.002*
Subjective Global Assessment: (mean ±SD)	2.02± .69		1.48± .50		Paired t test p< 0.001*
MRC dyspnea scale: (mean ±SD)	2.70±.68		1.99±.77		Paired t test p< 0.001*
COPD assessment test: (mean ±SD)	19.47±5.75		12.98±5.63		Paired t test p< 0.001*

Abbreviation: SD (standard deviation)/ ** significant/ SGA (Subjective global assessment)/ MRC dyspnea scale (modified research council dyspnea scale)/ HRQoL (health related quality of life)/CAT (COPD assessment test)

Discussion

Dietary counseling improved malnourished COPD patients' nutritional status, functional outcomes, and quality of life by increasing energy and protein intake, changing eating habits, preventing weight loss, promoting weight gain, and improving nutritional status, functional outcomes, and quality of life. The current study revealed after dietary counseling, there were improvement in nutritional status and decrease in malnutrition from three quarter (SGA B, SGAC) to nearly one-half percentage (SGA B, SGAC). Mean difference of CAT assessment test, MRC dyspnea showed improved score from pre to post intervention. Our findings are consistent with those of Nguyen *et al.*, who investigated the effectiveness of individualized nutritional counseling in treating malnourished outpatients with COPD in a low- & middle-income country (Vietnam). They found that planned nutrition advice had great effect in managing malnourished persons in underdeveloped nations without fortifying food or utilizing oral nutrition supplementation (Nguyen *et al.* 2020) ^[18].

Our results are also in line with those of Weekes *et al.*, who discovered nutrition counseling to be beneficial, however, the intervention group received both dietary advice and a supply of milk powder over a 6-month period. (Weekes *et al.* 2009) ^[23].

Yet our findings contradict those reported by Parsons *et al.*, 2017 ^[20] who compared the efficacy of ONS versus dietary advice in treating patients suffering from malnutrition in care homes in England. Those authors noted that after three months of the intervention, those who received dietary advice, did not show any improvement in nutritional status, inversely they reduced their energy and protein intake. This could be due to the fact that the dietary advice in that study was provided through a written resource that supported the consumption of energy-rich foods rather than being individually tailored. (Parsons *et al.* 2017) ^[17].

In a similar study done by Gupta *et al.* in India and Cochrane Review in the United Kingdom, no significant changes in nutritional status, lung function, or HRQOL were detected over time or between groups (intervention and control groups). (Gupta *et al.* 2014, Cochrane Review. 2004) ^[11, 6].

These previous findings might be attributed to the fact that the counseling messages did not mention why participants needed to improve their nutrient intake, or the adverse effects of reducing them and effect of weight loss on their health status. (Andersson *et al.* 2014, Rosenstock *et al.* 1988, Vermeeren. 2004, Efthimiou *et al.* 2004) ^[1, 21, 22, 8].

Also, Hallin *et al.* 2006 ^[12] did not found any improvement in nutritional status of their studied patients, they suggested

that chronic inflammation increase cytokine resulted in increased levels of leptin. These elevated leptin levels could result in a reduction in food intake and therefore poor response to nutritional support. "A cytokine–leptin theory" (Hallin *et al.* 2006) ^[12].

In our study there were significant relationship between malnourished patient and MRC dyspnea scale, CAT score and HRQOL, similarly Gunya *et al.* 2013 ^[10] in Turkey, demonstrated low pulmonary functions, MRC dyspnea scale among patients who were moderately and severely malnourished compared with well-nourished patients. Gupta B *et al.* 2010 ^[11] in India, described a significant relationship between poorer nourishment states and loss in pulmonary function test (PFT) results (especially in FEV1) in COPD patients. Yuceede *et al.* 2013 ^[24] in Turkey, stated that pulmonary functions were lower in the SGA-malnutrition group compared with the SGA-normal nutrition group. low body weight and long smoking history were associated with greater impairment in HRQoL score. Research has indicated that a four-point improvement in HRQoL score can contribute to a change in clinical outcomes. (Nguyen *et al.* 2019) ^[17].

Our participants had their HRQOL markedly impaired across all levels of malnutrition and COPD severity. Monsó *et al.* 1998 ^[16] in Germany, in their study on 47 COPD patients they found no association between HRQoL and nutritional status. Similarly, Monsó *et al.* 1998 ^[16] in Germany, in their study on 47 COPD patients they found no association between HRQoL and nutritional status while in a study by Cömert *et al.* 2011 ^[7] in Turkey, Gunya *et al.* 2013 ^[10] in Turkey, Nguyen *et al.* 2019 in Vietnam reported a significant correlation between HRQoL and nutritional status this may be due to variation in instrument used to measure nutritional status and HRQoL.

Declaration

Ethical Standards Disclosure: "This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving research study participants were approved by the [ethics committee of Tanta university], (approval number 33745/3/20). Formal consent was obtained from all subjects/patients.

Consent for publication: "None." As my manuscript does not contains any individual person's data in any form.

Data availability: The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Conflict of Interest: "None."

Financial Support/funding: "None."

Acknowledgements

[My Mother, Father, and Family that provided advice and support]

Authorship: [SHK: Conceptualization, methodology, data analysis and writing the original draft. RE: reviewed the findings and the original draft. AA: revision of the final manuscript for submission. NH: supervision and direction throughout the research, reviewing methodology, data curation, the original draft and final manuscript for

submission]. All authors read and approved the final manuscript.

References

- Andersson J, Hulander E, Rothenberg E, Iversen PO. Effect on body weight, quality of life and appetite following individualized, nutritional counselling to home-living elderly after rehabilitation–An open randomized trial. *The journal of nutrition, health & aging.* 2017 Jul;21(7):811-818. <https://doi.org/10.1007/s12603-016-0825-8>
- Baldwin C, Weekes CE. Dietary advice with or without oral nutritional supplements for disease-related malnutrition in adults. *Cochrane database of systematic reviews,* 2011, 9. <https://doi.org/10.1002/14651858.CD002008.pub4>
- British Association for Parenteral and Enteral Nutrition. BDA nutrition and dietetic care process; c2016. <https://www.bapen.org.uk/nutrition-support/assessment-and-planning/nutritional-assessment>. Accessed January 20, 2021.
- Canadian Malnutrition Task Force; c2021. available at <https://nutritioncareinCanada.ca/resources-and-tools/hospital-care-inpac/assessment-sga?lang=en>. Accessed on January 20, 2021.
- CAT Development Steering Group. COPD assessment test–healthcare professional user guide; c2018. 2014-05], available at <http://www.catestonline.org/images/UserGuides/CATHCPUser%20guideEn.pdf>.
- Cochrane WJ, Afolabi OA. Investigation into the nutritional status, dietary intake and smoking habits of patients with chronic obstructive pulmonary disease. *Journal of human nutrition and dietetics.* 2004 Feb;17(1):3-11. <https://doi.org/10.1046/j.1365-277x.2003.00478.x>
- Cömert S, Erk M, Müsellim B, *et al.* Kronik obstrüktif akciğer hastalığında yaşam kalitesini etkileyen fonksiyonel ve antropometrik parametreler. *İzmir Göğüs Hastanesi Dergisi.* 2011;25:89-100.
- Efthimiou J, Fleming J, Gomes C, Spiro SG. The effect of supplementary oral nutrition in poorly nourished patients with chronic obstructive pulmonary disease. *American Review of Respiratory Disease.* 1988;137(5):1075-1082. <https://doi.org/10.1164/ajrccm/137.5.1075>
- Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease goldcopd; c2017. [org/wp-content/uploads/2017/11/GOLD-2018-v6.0-FINAL-revised-20-Nov_WMS.pdf](http://www.goldcopd.org/wp-content/uploads/2017/11/GOLD-2018-v6.0-FINAL-revised-20-Nov_WMS.pdf). Accessed on January 20, 2021.
- Günay E, Kaymaz D, Selçuk NT, Ergün P, Sengül F, Demir N. Effect of nutritional status in individuals with chronic obstructive pulmonary disease undergoing pulmonary rehabilitation. *Respirology;* c2013. Nov;18(8):1217-22. <https://doi.org/10.1111/resp.12133>
- Gupta B, Kant S, Mishra R. Effect of nutrition education intervention on symptomatic markers of Indian patients with chronic obstructive pulmonary disease. *J Nutr Health Food Eng.* 2014;1(3):105-14. <https://doi.org/10.15406/jnhfe.2014.01.00016>
- Hallin R, Koivisto-Hursti UK, Lindberg E, Janson C.

- Nutritional status, dietary energy intake and the risk of exacerbations in patients with chronic obstructive pulmonary disease (COPD). *Respiratory medicine*. 2006 Mar 1;100(3):561-567.
<https://doi.org/10.1016/j.rmed.2005.05.020>
13. Hoong JM, Ferguson M, Hukins C, Collins PF. Economic and operational burden associated with malnutrition in chronic obstructive pulmonary disease. *Clinical Nutrition*. 2017 Aug 1;36(4):1105-1109.
<https://doi.org/10.1016/j.clnu.2016.07.008>
 14. Ingadottir AR, Beck AM, Baldwin C, Weekes CE, Geirsdottir OG, Ramel A, *et al.* Oral nutrition supplements and between-meal snacks for nutrition therapy in patients with COPD identified as at nutritional risk: A randomised feasibility trial. *BMJ open respiratory research*. 2019 Jan 1;6(1).
<https://doi.org/10.1016/j.clnu.2018.06.1346>
 15. Medical Research Council. Questionnaire on respiratory symptoms:1986 instructions to interviewers. London: MRC Committee on Environmental and Occupational Health; c1986.
 16. Monsó E, Fiz JM, Izquierdo J, *et al.* Quality of life in severe chronic obstructive pulmonary disease: correlation with lung and muscle function. *Respir. Med*. 1998;92(2):221-227.
 17. Nguyen HT, Collins PF, Pavey TG, Nguyen NV, Pham TD, Gallegos DL. Nutritional status, dietary intake, and health-related quality of life in outpatients with COPD. *International journal of chronic obstructive pulmonary disease*. 2019;14:215.
<https://doi.org/10.2147/copd.s181322> Nguyen HT,
 18. Pavey TG, Collins PF, Nguyen NV, Pham TD, Gallegos D. Effectiveness of tailored dietary counseling in treating malnourished outpatients with chronic obstructive pulmonary disease: A randomized controlled trial. *Journal of the Academy of Nutrition and Dietetics*. 2020 May 1;120(5):778-791.
<https://doi.org/10.1016/j.jand.2019.09.013>
 19. Nguyen Viet N, Yunus F, Nguyen Thi Phuong A, Dao Bich V, Damayanti T, Wiyono WH, *et al.* The prevalence and patient characteristics of chronic obstructive pulmonary disease in non-smokers in Vietnam and Indonesia: An observational survey. *Respirology*. 2015 May;20(4):602-611.
<https://doi.org/10.1111/resp.12507>
 20. Parsons EL, Stratton RJ, Cawood AL, Smith TR, Elia M. Oral nutritional supplements in a randomized trial are more effective than dietary advice at improving quality of life in malnourished care home residents. *Clinical Nutrition*. 2017 Feb 1;36(1):134-142.
<https://doi.org/10.1016/j.clnu.2016.01.002>
 21. Rosenstock IM, Strecher VJ, Becker MH. Social learning theory and the Health Belief Model. *Health Educ Q*. Summer. 1988;15(2):175-83. Doi: 10.1177/109019818801500203. PMID: 3378902.
 22. Vermeeren MA, Wouters EF, Geraerts-Keeris AJ, Schols AM. Nutritional support in patients with chronic obstructive pulmonary disease during hospitalization for an acute exacerbation; a randomized controlled feasibility trial. *Clinical nutrition*. 2004 Oct 1;23(5):1184-192.
<https://doi.org/10.1016/j.clnu.2004.03.008>
 23. Weekes CE, Emery PW, Elia M. Dietary counselling and food fortification in stable COPD: a randomized trial. *Thorax*. 2009 Apr 1;64(4):326-331.
<https://doi.org/10.1136/thx.2008.097352>
 24. Yuceege MB, Salman SO, Duru SM, Saygideger YM, Sonmez Z, Ardic S. The Evaluation of nutrition in male COPD patients using subjective global assessment and mini nutritional assessment. *Int J Intern Med*. 2013;2(1):1-5.

How to Cite This Article

Koabar SMM, El-Salamy RM, Abd El-Zaher AH, Hassan NM. Effect of dietary counseling on patients with chronic obstructive pulmonary disease admitted to a tertiary care hospital, Middle Delta, Egypt. *International Journal of Advanced Community Medicine*. 2023;6(1):01-06.

Creative Commons (CC) License

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.